







# Methylation







#### References

- Duan Z. et al. (2012) A genome-wide 3C-method for characterizing the three-dimensional architectures of genomes. Methods 58: 277-288
- 4-C Zhao Z. et al. (2006) Circular chromosome conformation capture (4C) uncovers extensive networks of epigenetically regulated intra- and interchromosomal interactions. Nat Genet 38: 1341-1347
- 5-C Dostie J. et al. (2007) Mapping networks of physical interactions between genomic elements using 5C technology. Nat Protoc 2: 988-1002 Buenrostro J. D. et al. (2013) Transposition of native chromatin for fast and sensitive epig-ATAC-Seq CLASH
- enomic profiling of open chromatin, DNA-binding proteins and nucleosome position. Nat Methods CLIP-Seq
- Tani H. et al. (2012) Genome-wide determination of RNA stability reveals hundreds of BRIC-seq short-lived noncoding transcripts in mammals. Genome Res 22: 947-956
- Paulsen M. T. et al. (2013) Coordinated regulation of synthesis and stability of RNA during the Bru-Sea acute TNF-induced proinflammatory response. Proc Natl Acad Sci U S A 110: 2240-2245
- BruChase-Seg Paulsen M, T, et al. (2013) Coordinated regulation of synthesis and stability of RNA during the acute TNF-induced proinflammatory response. Proc Natl Acad Sci U S A 110: 2240-2245 Feil R., Charlton J., Bird A. P., Walter J. and Reik W. (1994) Methylation analysis on individual BS-Seq
- chromosomes: improved protocol for bisulphite genomic sequencing. Nucleic Acids Res 22: 695-696
- Cap-Seq Illingworth R. S. et al. (2010) Orphan CpG islands identify numerous conserved promoters in
- Cell 129: 823-837

DNase-Seq

- ChIRP-Sea Chu C. et al. (2011) Genomic maps of long noncoding RNA occupancy reveal principles of GMUCT 1.0 Gregory B. D. et al. (2008) A link between RNA metabolism and silencing affecting Arabidopsis MDA RNA-chromatin interactions. Mol Cell 44: 667-678 CIP-TAP Gu W. et al. (2012) CapSeq and CIP-TAP identify Pol II start sites and reveal capped small GMUCT 2.0 RNAs as C. elegans piRNA precursors. Cell 151: 1488-1500 Cir-Seq Acevedo A. et al. (2014) Mutational and fitness landscapes of an RNA virus revealed through GRO-Seq population sequencing. Nature 505: 686-690
  - Kudla G. et al. (2011) Cross-linking, ligation, and sequencing of hybrids reveals RNA-RNA Hi-C interactions in yeast. Proc Natl Acad Sci U S A 108: 10010-10015
- See RIP-Sea Digital RNA Shiroguchi K. et al. (2012) Digital RNA sequencing minimizes sequence-dependent bias and amplification noise with optimized single-molecule barcodes. Proc Natl Acad Sci U S A 109: HITS-CLIP
  - 1347-1352 Boyle A. P. et al. (2008) High-Resolution Mapping and Characterization of Open Chromatin ICE across the Genome. Cell 132: 311-322
- Hesselberth J. R. et al. (2009) Global mapping of protein-DNA interactions in vivo by digital iCLIP DNaseI-Seq genomic footprinting. Nat Methods 6: 283-289 DP-Seq
  - Bhargava V. et al. (2013) Quantitative transcriptomics using designed primer-based amplifica- Ig-Seq tion. Sci Rep 3: 1740
- development. Dev Cell 14: 854-866 scripts in eukaryotes-GMUCT 2.0. Methods 67: 64-73 Core L. J. et al. (2008) Nascent RNA sequencing reveals widespread pausing and divergent MeDIP-Seq initiation at human promoters. Science 322: 1845-1848 Lieberman-Aiden E. et al. (2009) Comprehensive mapping of long-range interactions reveals MeRIP-Seq
- folding principles of the human genome. Science 326: 289-293 Histone methylation Barski A. et al. (2007) High-resolution profiling of histone methylations in the human genome. MethylCap-Seq See MDBCap-Seq Cell 129: 823-837
  - Chi S. W. et al. (2009) Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. Nature 460: 479-486
  - Sakurai M. et al. (2010) Inosine cyanoethylation identifies A-to-I RNA editing sites in the human transcriptome. Nat Chem Biol 6: 733-740 Konig J. et al. (2010) iCLIP reveals the function of hnRNP particles in splicing at individual
  - nucleotide resolution. Nat Struct Mol Biol 17: 909-915 Vollmers C. et al. (2013) Genetic measurement of memory B-cell recall using antibody repertoire sequencing. Proc Natl Acad Sci U S A 110: 13463-13468
- Dean F. B. et al. (2001) Rapid amplification of plasmid and phage DNA using Phi 29 DNA poly- PEAT merase and multiply-primed rolling circle amplification. Genome Res 11: 1095-1099 Willmann M. R. et al. (2014) Improved genome-wide mapping of uncapped and cleaved tran- MDBCap-Seq de Assis S. et al. (2012) High-fat or ethinyl-oestradiol intake during pregnancy increases Quartz-Seq mammary cancer risk in several generations of offspring. Nat Commun 3: 1053
  - Down T. A. et al. (2008) A Bayesian deconvolution strategy for immunoprecipitation-based DNA methylome analysis. Nat Biotechnol 26: 779-785
  - Meyer K. D. et al. (2012) Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. Cell 149: 1635-1646

  - See MAINE-Seq MNase-Seg NFT-Sea
    - Churchman L. S. et al. (2011) Nascent transcript sequencing visualizes transcription at nucleotide resolution. Nature 469: 368-373 Han H. et al. (2011) DNA methylation directly silences genes with non-CpG island promoters NOME-Sea
    - and establishes a nucleosome occupied promoter. Hum Mol Genet 20: 4299-4310 Os-Sea Myllykangas S. et al. (2011) Efficient targeted resequencing of human germline and cancer
    - genomes by oligonucleotide-selective sequencing. Nat Biotechnol 29: 1024-1027 oxBS-Seq Booth M. J. et al. (2012) Quantitative sequencing of 5-methylcytosine and 5-hydroxymethyl
      - cytosine at single-base resolution. Science 336: 934-937
- Ni T. et al. (2010) A paired-end sequencing strategy to map the complex landscape of tran- SMORE-seq scription initiation. Nat Methods 7: 521-527
- Sasagawa Y. et al. (2013) Quartz-Seq: a highly reproducible and sensitive single-cell RNA sequencing method, reveals non-genetic gene-expression heterogeneity. Genome Biol 14: STRT

D<sup>-</sup> P<sup>-</sup> O S

RAD

RIP-Sea

SHAPE-Sea

- Baird N. A. et al. (2008) Rapid SNP discovery and genetic mapping using sequenced RAD Structure-Seq markers. PLoS ONE 3: e3376
- Paired-end RAD-Seq Willing E. M. et al. (2011) Paired-end RAD-seq for de novo assembly and marker design without available reference. Bioinformatics 27: 2187-2193
- Baillie J. K. et al. (2011) Somatic retrotransposition alters the genetic landscape of the human RC-Seq brain. Nature 479: 534-537 Ribo-Seq
  - Ingolia N. T. et al. (2009) Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. Science 324: 218-223 Zhao J. et al. (2010) Genome-wide identification of polycomb-associated RNAs by RIP-seq. Mol
- Cell 40: 939-953 Xi Y. et al. (2012) RRBSMAP: a fast, accurate and user-friendly alignment tool for reduced RRBS-Seq
  - representation bisulfite sequencing. Bioinformatics 28: 430-432 Lucks J. B. et al. (2011) Multiplexed RNA structure characterization with selective 2'-hydroxyl
- Park D. et al. (2014) Simultaneous mapping of transcript ends at single-nucleotide resolution and identification of widespread promoter-associated non-coding RNA governed by TATA elements. Nucleic Acids Res
- Islam S. et al. (2011) Characterization of the single-cell transcriptional landscape by highly multiplex RNA-seq. Genome Res 21: 1160-1167
- Ding Y. et al. (2014) In vivo genome-wide profiling of RNA secondary structure reveals novel regulatory features. Nature 505: 696-700
  - Schapire A. L. et al. (2013) Construction of Specific Parallel Amplification of RNA Ends (SPARE) libraries for the systematic identification of plant microRNA processing intermediates. Methods 64:283-291
- TAB-Seq Yu M. et al. (2012) Base-resolution analysis of 5-hydroxymethylcytosine in the Mammalian genome. Cell 149: 1368-1380
- TC-Seq Klein IA (2011) Translocation-capture sequencing reveals the extent and nature of chromosomal rearrangements in B lymphocytes. Cell 147:95-106.
- TCR Chain Paring Turchaninova M. A. et al. (2013) Pairing of T-cell receptor chains via emulsion PCR. Eur J Immunol 43: 2507-2515
  - Pelechano V. et al. (2013) Extensive transcriptional heterogeneity revealed by isoform profiling. Nature 497: 127-131
- van Opijnen T. et al. (2013) Transposon insertion sequencing: a new tool for systems-level



- Hashimshony T. et al. (2012) CEL-Seq: single-cell RNA-Seq by multiplexed linear amplification. Cell Rep 2: 666-673
- Li G. et al. (2010) ChIA-PET tool for comprehensive chromatin interaction analysis with Chia-PET paired-end tag sequencing. Genome Biol 11: R22
- ChIP-exo Yen K. et al. (2013) SWR-C and INO80 Chromatin Remodelers Recognize Nucleosome-free Regions Near +1 Nucleosomes. Cell 154: 1246-1256
- ChIP-Seq Barski A. et al. (2007) High-resolution profiling of histone methylations in the human genome.
- Zheng Q. et al. (2010) Genome-wide double-stranded RNA sequencing reveals the functional IMS-MDA dsRNA-Seq significance of base-paired RNAs in Arabidopsis. PLoS Genet 6: e1001141 Duplex-Seq Schmitt M. W. et al. (2012) Detection of ultra-rare mutations by next-generation sequencing. INSeq Proc Natl Acad Sci U S A 109: 14508-14513 FAIRE-Seq Gaulton K. J. et al. (2010) A map of open chromatin in human pancreatic islets. Nat Genet 42: MAINE-Seq
- 255-259 Underwood J. G. et al. (2010) FragSeq: transcriptome-wide RNA structure probing using MALBAC Frag-Seq
  - high-throughput sequencing. Nat Methods 7: 995-1001
- Seth-Smith H. M. et al. (2013) Generating whole bacterial genome sequences of low-abun-PAR-CLTP dance species from complex samples with IMS-MDA. Nat Protoc 8: 2404-2412
- van Opijnen T. et al. (2013) Transposon insertion sequencing: a new tool for systems-level PARE-Sea analysis of microorganisms. Nat Rev Microbiol 11: 435-442 Ponts N. et al. (2010) Nucleosome landscape and control of transcription in the human malaria
- parasite. Genome Res 20: 228-238
- DVDC Zong C. et al. (2012) Genome-wide detection of single-nucleotide and copy-number variations DR A1 of a single human cell. Science 338: 1622-1626
- Hafner M. et al. (2010) Transcriptome-wide identification of RNA-binding protein and microR-NA target sites by PAR-CLIP. Cell 141: 129-141
- Smart-Seg German M. A. et al. (2008) Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. Nat Biotechnol 26: 941-946
- Smart-seq2 Wan Y. et al. (2013) Genome-wide mapping of RNA structure using nuclease digestion and high-throughput sequencing. Nat Protoc 8: 849-869 smMIP
  - Miura F. et al. (2012) Amplification-free whole-genome bisulfite sequencing by post-bisulfite adaptor tagging. Nucleic Acids Res 40: e136
- acylation analyzed by primer extension sequencing (SHAPE-Seq). Proc Natl Acad Sci U S A TN-Seq 108: 11063-11068 Ramskold D. et al. (2012) Full-length mRNA-Seq from single-cell levels of RNA and individual TRAP-Seq
- new levels of control. Mol Syst Biol 6: 419 circulating tumor cells. Nat Biotechnol 30: 777-782 Picelli S. et al. (2013) Smart-seq2 for sensitive full-length transcriptome profiling in single T-WGBS

TIF-Seq

- cells. Nat Methods 10: 1096-1098v Hiatt JB (2013) Single molecule molecular inversion probes for targeted, high-accuracy detec- UMI Method
- tion of low-frequency variation. Genome Res 23:843-54.
- analysis of microorganisms. Nat Rev Microbiol 11: 435-442 Jiao Y. et al. (2010) Cell-type specific analysis of translating RNAs in developing flowers reveals
- Wang Q. et al. (2013) Tagmentation-based whole-genome bisulfite sequencing. Nat Protoc 8: 2022-2032
- Kivioja T. et al. (2012) Counting absolute numbers of molecules using unique molecular identifiers. Nat Methods 9: 72-74





### ~10kb Sheared genomic DNA End repair Adapter ligation \_ Generate clonal pools Amplify \_\_\_\_ Tagmentation ~600bp Add indices Pool and purify P5 \_\_\_\_\_ P7 Prepared fragments

Illumina, Inc. • 5200 Illumina Way, San Diego, CA 92122 USA • 1.800.809.4566 toll-free • 1.858.202.4566 tel • techsupport@illumina.com • illumina.com

## FOR RESEARCH USE ONLY

## © 2014 Illumina, Inc. All rights reserved.

Illumina, IlluminaDx, BaseSpace, BeadArray, BeadXpress, cBot, CSPro, DASL, DesignStudio, Eco, GAIIx, Genetic Energy, Genome Analyzer, GenomeStudio, GoldenGate, HiSeq, Infinium, iSelect, MiSeq, Nextera, NextSeq, NuPCR, SeqMonitor, Solexa, TruSeq, TruSight, VeraCode, the pumpkin orange color, and the Genetic Energy streaming bases design are trademarks of Illumina, Inc. All other brands and names contained herein are the property of their respective owners.

